

# Full Reference List of Blood Sugar Clinical Studies for Supergut

## Blood Sugar Management: Resistant Starch Studies (85)

1. [Effects of dietary propionate on carbohydrate and lipid metabolism in healthy volunteers.](#)  
Venter CS, Vorster HH, Cummings JH.  
*Am J Gastroenterol.* 1990 May;85(5):549-53.  
PMID: 2159696 Clinical Trial.
2. [Intestinal transport and fermentation of resistant starch evaluated by the hydrogen breath test.](#)  
Olesen M, Rumessen JJ, Gudmand-Høyer E.  
*Eur J Clin Nutr.* 1994 Oct;48(10):692-701.  
PMID: 7835324 Clinical Trial.
3. [Nutrients excreted in ileostomy effluents after consumption of mixed diets with beans or potatoes. II. Starch, dietary fibre and sugars.](#)  
Schweizer TF, Andersson H, Langkilde AM, Reimann S, Torsdottir I.  
*Eur J Clin Nutr.* 1990 Aug;44(8):567-75.  
PMID: 2170104 Clinical Trial.
4. [Resistant starch: the effect on postprandial glycemia, hormonal response, and satiety.](#)  
Raben A, Tagliabue A, Christensen NJ, Madsen J, Holst JJ, Astrup A.  
*Am J Clin Nutr.* 1994 Oct;60(4):544-51. doi: 10.1093/ajcn/60.4.544.  
PMID: 8092089 Clinical Trial.
5. [Physiological effects of resistant starches on fecal bulk, short chain fatty acids, blood lipids and glycemic index.](#)  
Jenkins DJ, Vuksan V, Kendall CW, Würsch P, Jeffcoat R, Waring S, Mehling CC, Vidgen E, Augustin LS, Wong E.  
*J Am Coll Nutr.* 1998 Dec;17(6):609-16. doi: 10.1080/07315724.1998.10718810.  
PMID: 9853541 Clinical Trial.
6. [Acute effect of fructose on postprandial lipaemia in diabetic and non-diabetic subjects.](#)  
Abraha A, Humphreys SM, Clark ML, Matthews DR, Frayn KN.  
*Br J Nutr.* 1998 Aug;80(2):169-75.  
PMID: 9828758 Clinical Trial.
7. [Glycaemic and insulinaemic responses to natural foods, frozen foods and their laboratory equivalents.](#)  
Kanan W, Bijlani RL, Sachdeva U, Mahapatra SC, Shah P, Karmarkar MG.  
*Indian J Physiol Pharmacol.* 1998 Jan;42(1):81-9.  
PMID: 9513797 Clinical Trial.
8. [Effect of high-amyllose starch and oat bran on metabolic variables and bowel function in subjects with hypertriglyceridemia.](#)  
Noakes M, Clifton PM, Nestel PJ, Le Leu R, McIntosh G.  
*Am J Clin Nutr.* 1996 Dec;64(6):944-51. doi: 10.1093/ajcn/64.6.944.  
PMID: 8942421 Clinical Trial.
9. [Neither raw nor retrograded resistant starch lowers fasting serum cholesterol concentrations in healthy normolipidemic subjects.](#)

- Heijnen ML, van Amelsvoort JM, Deurenberg P, Beynen AC.  
Am J Clin Nutr. 1996 Sep;64(3):312-8. doi: 10.1093/ajcn/64.3.312.  
PMID: 8780339 Clinical Trial.
10. [The influence of starch structure on glycogen resynthesis and subsequent cycling performance.](#)  
Jozsi AC, Trappe TA, Starling RD, Goodpaster B, Trappe SW, Fink WJ, Costill DL.  
Int J Sports Med. 1996 Jul;17(5):373-8. doi: 10.1055/s-2007-972863.  
PMID: 8858410 Clinical Trial.
11. [The effects of pre-exercise starch ingestion on endurance performance.](#)  
Goodpaster BH, Costill DL, Fink WJ, Trappe TA, Jozsi AC, Starling RD, Trappe SW.  
Int J Sports Med. 1996 Jul;17(5):366-72. doi: 10.1055/s-2007-972862.  
PMID: 8858409 Clinical Trial.
12. [Dietary fibre, resistant starch and in vitro starch digestibility of cereal meals. Glycaemic and insulinaemic responses in NIDDM patients.](#)  
Lintas C, Cappelloni M, Bonmassar L, Clementi A, Del Toma E, Ceccarelli G.  
Eur J Clin Nutr. 1995 Oct;49 Suppl 3:S264-7.  
PMID: 8549543 Clinical Trial. No abstract available.
13. [Sourdough fermentation or addition of organic acids or corresponding salts to bread improves nutritional properties of starch in healthy humans.](#)  
Liljeberg HG, Lönnér CH, Björck IM.  
J Nutr. 1995 Jun;125(6):1503-11. doi: 10.1093/jn/125.6.1503.  
PMID: 7782904 Clinical Trial.
14. [Replacement of digestible by resistant starch lowers diet-induced thermogenesis in healthy men.](#)  
Heijnen ML, Deurenberg P, van Amelsvoort JM, Beynen AC.  
Br J Nutr. 1995 Mar;73(3):423-32. doi: 10.1079/bjn19950044.  
PMID: 7766565 Clinical Trial.
15. [\[Effects of resistant starch on insulin resistance of type 2 diabetes mellitus patients\].](#)  
Zhang WQ, Wang HW, Zhang YM, Yang YX.  
Zhonghua Yu Fang Yi Xue Za Zhi. 2007 Mar;41(2):101-4.  
PMID: 17605234 Clinical Trial. Chinese.
16. [Insulin-sensitizing effects of dietary resistant starch and effects on skeletal muscle and adipose tissue metabolism.](#)  
Robertson MD, Bickerton AS, Dennis AL, Vidal H, Frayn KN.  
Am J Clin Nutr. 2005 Sep;82(3):559-67. doi: 10.1093/ajcn.82.3.559.  
PMID: 16155268 Clinical Trial.
17. [Effect of bread containing resistant starch on postprandial blood glucose levels in humans.](#)  
Yamada Y, Hosoya S, Nishimura S, Tanaka T, Kajimoto Y, Nishimura A, Kajimoto O.  
Biosci Biotechnol Biochem. 2005 Mar;69(3):559-66. doi: 10.1271/bbb.69.559.  
PMID: 15784985 Clinical Trial.
18. [Resistant starch supplementation influences blood lipid concentrations and glucose control in overweight subjects.](#)  
Park OJ, Kang NE, Chang MJ, Kim WK.  
J Nutr Sci Vitaminol (Tokyo). 2004 Apr;50(2):93-9.  
PMID: 15242012 Clinical Trial.
19. [Prior short-term consumption of resistant starch enhances postprandial insulin sensitivity in healthy subjects.](#)  
Robertson MD, Currie JM, Morgan LM, Jewell DP, Frayn KN.  
Diabetologia. 2003 May;46(5):659-65. doi: 10.1007/s00125-003-1081-0. Epub 2003 Apr 24.  
PMID: 12712245 Clinical Trial.

20. [Effect of the glycemic index and content of indigestible carbohydrates of cereal-based breakfast meals on glucose tolerance at lunch in healthy subjects.](#)  
Liljeberg HG, Akerberg AK, Björck IM.  
*Am J Clin Nutr.* 1999 Apr;69(4):647-55. doi: 10.1093/ajcn/69.4.647.  
PMID: 10197565 Clinical Trial.
21. [Resistant starch and pullulan reduce postprandial glucose, insulin, and GLP-1, but have no effect on satiety in healthy humans.](#)  
Klosterbuer AS, Thomas W, Slavin JL.  
*J Agric Food Chem.* 2012 Dec 5;60(48):11928-34. doi: 10.1021/jf303083r. Epub 2012 Nov 20.  
PMID: 23136915 Clinical Trial.
22. [Dietary treatment with rice containing resistant starch improves markers of endothelial function with reduction of postprandial blood glucose and oxidative stress in patients with prediabetes or newly diagnosed type 2 diabetes.](#)  
Kwak JH, Paik JK, Kim HI, Kim OY, Shin DY, Kim HJ, Lee JH, Lee JH.  
*Atherosclerosis.* 2012 Oct;224(2):457-64. doi: 10.1016/j.atherosclerosis.2012.08.003. Epub 2012 Aug 27.  
PMID: 22954674 Clinical Trial.
23. [The effect of yellow pea protein and fibre on short-term food intake, subjective appetite and glycaemic response in healthy young men.](#)  
Smith CE, Molland RC, Luhovyy BL, Anderson GH.  
*Br J Nutr.* 2012 Aug;108 Suppl 1:S74-80. doi: 10.1017/S0007114512000700.  
PMID: 22916818 Clinical Trial.
24. [Dietary fibre improves first-phase insulin secretion in overweight individuals.](#)  
Bodinham CL, Smith L, Wright J, Frost GS, Robertson MD.  
*PLoS One.* 2012;7(7):e40834. doi: 10.1371/journal.pone.0040834. Epub 2012 Jul 16.  
PMID: 22815837 **Free PMC article.** Clinical Trial.
25. [African americans may have to consume more than 12 grams a day of resistant starch to lower their risk for type 2 diabetes.](#)  
Penn-Marshall M, Holtzman GI, Barbeau WE.  
*J Med Food.* 2010 Aug;13(4):999-1004. doi: 10.1089/jmf.2009.0195.  
PMID: 20482275 Clinical Trial.
26. [Resistant starch improves insulin sensitivity in metabolic syndrome.](#)  
Johnston KL, Thomas EL, Bell JD, Frost GS, Robertson MD.  
*Diabet Med.* 2010 Apr;27(4):391-7. doi: 10.1111/j.1464-5491.2010.02923.x.  
PMID: 20536509 Clinical Trial.
27. [Relation between estimates of cornstarch digestibility by the Englyst in vitro method and glycemic response, subjective appetite, and short-term food intake in young men.](#)  
Anderson GH, Cho CE, Akhavan T, Molland RC, Luhovyy BL, Finocchiaro ET.  
*Am J Clin Nutr.* 2010 Apr;91(4):932-9. doi: 10.3945/ajcn.2009.28443. Epub 2010 Feb 17.  
PMID: 20164321 Clinical Trial.
28. [Postprandial glycaemic and insulinaemic responses to GM-resistant starch-enriched rice and the production of fermentation-related H2 in healthy Chinese adults.](#)  
Li M, Piao JH, Tian Y, Li WD, Li KJ, Yang XG.  
*Br J Nutr.* 2010 Apr;103(7):1029-34. doi: 10.1017/S0007114509992820. Epub 2009 Nov 24.  
PMID: 19930763 Clinical Trial.
29. [Dietary amylose and amylopectin ratio and resistant starch content affects plasma glucose, lactic acid, hormone levels and protein synthesis in splanchnic tissues.](#)  
Deng J, Wu X, Bin S, Li TJ, Huang R, Liu Z, Liu Y, Ruan Z, Deng Z, Hou Y, Yin YL.

- J Anim Physiol Anim Nutr (Berl). 2010 Apr;94(2):220-6. doi: 10.1111/j.1439-0396.2008.00902.x. Epub 2009 Jan 13.  
PMID: 19175452 Clinical Trial.
30. [Acute ingestion of resistant starch reduces food intake in healthy adults.](#)  
Bodinham CL, Frost GS, Robertson MD.  
Br J Nutr. 2010 Mar;103(6):917-22. doi: 10.1017/S0007114509992534. Epub 2009 Oct 27.  
PMID: 19857367 Clinical Trial.
31. [The effects of whole grain high-amylose maize flour as a source of resistant starch on blood glucose, satiety, and food intake in young men.](#)  
Luhovyy BL, Mollard RC, Yurchenko S, Nunez MF, Berengut S, Liu TT, Smith CE, Pelkman CL, Anderson GH.  
J Food Sci. 2014 Dec;79(12):H2550-6. doi: 10.1111/1750-3841.12690. Epub 2014 Nov 11.  
PMID: 25388622 Clinical Trial.
32. [Effects of resistant starch on behaviour, satiety-related hormones and metabolites in growing pigs.](#)  
Souza da Silva C, Haenen D, Koopmans SJ, Hooiveld GJ, Bosch G, Bolhuis JE, Kemp B, Müller M, Gerrits WJ.  
Animal. 2014 Sep;8(9):1402-11. doi: 10.1017/S1751731114001116. Epub 2014 May 20.  
PMID: 24845880 Clinical Trial.
33. [Resistant starch type 4-enriched diet lowered blood cholesterol and improved body composition in a double blind controlled cross-over intervention.](#)  
Nichenametla SN, Weidauer LA, Wey HE, Beare TM, Specker BL, Dey M.  
Mol Nutr Food Res. 2014 Jun;58(6):1365-9. doi: 10.1002/mnfr.201300829. Epub 2014 Jan 30.  
PMID: 24478107 **Free PMC article.** Clinical Trial.
34. [Grain sorghum muffin reduces glucose and insulin responses in men.](#)  
Poquette NM, Gu X, Lee SO.  
Food Funct. 2014 May;5(5):894-9. doi: 10.1039/c3fo60432b.  
PMID: 24608948 Clinical Trial.
35. [Endogenous plasma glucagon-like peptide-1 following acute dietary fibre consumption.](#)  
Bodinham CL, Al-Mana NM, Smith L, Robertson MD.  
Br J Nutr. 2013 Oct;110(8):1429-33. doi: 10.1017/S0007114513000731. Epub 2013 Mar 18.  
PMID: 23507477 Clinical Trial.
36. [Postprandial glucose, insulin and gastrointestinal hormones in healthy and diabetic subjects fed a fructose-free and resistant starch type IV-enriched enteral formula.](#)  
García-Rodríguez CE, Mesa MD, Olza J, Buccianti G, Pérez M, Moreno-Torres R, Pérez de la Cruz A, Gil A.  
Eur J Nutr. 2013 Sep;52(6):1569-78. doi: 10.1007/s00394-012-0462-x. Epub 2012 Nov 23.  
PMID: 23179201 Clinical Trial.
37. [Impact of short term consumption of diets high in either non-starch polysaccharides or resistant starch in comparison with moderate weight loss on indices of insulin sensitivity in subjects with metabolic syndrome.](#)  
Lobley GE, Holtrop G, Bremner DM, Calder AG, Milne E, Johnstone AM.  
Nutrients. 2013 Jun 10;5(6):2144-72. doi: 10.3390/nu5062144.  
PMID: 23752495 **Free PMC article.** Clinical Trial.
38. [Low dose chromium-polynicotinate or policosanol is effective in hypercholesterolemic children only in combination with glucomannan.](#)  
Martino F, Puddu PE, Pannarale G, Colantoni C, Martino E, Niglio T, Zanoni C, Barillà F.  
Atherosclerosis. 2013 May;228(1):198-202. doi: 10.1016/j.atherosclerosis.2013.02.005.  
Epub 2013 Feb 13.

- PMID: 23453352 Clinical Trial.
39. [On the possibility to affect the course of glycaemia, insulinaemia, and perceived hunger/satiety to bread meals in healthy volunteers.](#)  
Ekström LM, Björck IM, Östman EM.  
Food Funct. 2013 Apr 25;4(4):522-9. doi: 10.1039/c2fo30251a. Epub 2013 Jan 21.  
PMID: 23334658 Clinical Trial.
40. [Effects of indigestible carbohydrates in barley on glucose metabolism, appetite and voluntary food intake over 16 h in healthy adults.](#)  
Johansson EV, Nilsson AC, Östman EM, Björck IM.  
Nutr J. 2013 Apr 11;12:46. doi: 10.1186/1475-2891-12-46.  
PMID: 23577719 **Free PMC article.** Clinical Trial.
41. [The Therapeutic Potential of Resistant Starch in Modulation of Insulin Resistance, Endotoxemia, Oxidative Stress and Antioxidant Biomarkers in Women with Type 2 Diabetes: A Randomized Controlled Clinical Trial.](#)  
Karimi P, Farhangi MA, Sarmadi B, Gargari BP, Zare Javid A, Pouraghaei M, Dehghan P.  
Ann Nutr Metab. 2016;68(2):85-93. doi: 10.1159/000441683. Epub 2015 Dec 12.  
PMID: 26655398 Clinical Trial.
42. [Is there any place for resistant starch, as alimentary prebiotic, for patients with type 2 diabetes?](#)  
Gargari BP, Namazi N, Khalili M, Sarmadi B, Jafarabadi MA, Dehghan P.  
Complement Ther Med. 2015 Dec;23(6):810-5. doi: 10.1016/j.ctim.2015.09.005. Epub 2015 Sep 16.  
PMID: 26645521 Clinical Trial.
43. [Glycemic Response and Fermentation of Crystalline Short Linear α-Glucans from Debranched Waxy Maize Starch.](#)  
Brewer LR, Weber C, Haub M, Cai L, Shi YC.  
J Agric Food Chem. 2015 Nov 4;63(43):9528-35. doi: 10.1021/acs.jafc.5b03632. Epub 2015 Oct 26.  
PMID: 26447350 Clinical Trial.
44. [Manipulation of starch bioaccessibility in wheat endosperm to regulate starch digestion, postprandial glycemia, insulinemia, and gut hormone responses: a randomized controlled trial in healthy ileostomy participants.](#)  
Edwards CH, Grundy MM, Grassby T, Vasilopoulou D, Frost GS, Butterworth PJ, Berry SE, Sanderson J, Ellis PR.  
Am J Clin Nutr. 2015 Oct;102(4):791-800. doi: 10.3945/ajcn.114.106203. Epub 2015 Sep 2.  
PMID: 26333512 **Free PMC article.** Clinical Trial.
45. [Increased gut hormones and insulin sensitivity index following a 3-d intervention with a barley kernel-based product: a randomised cross-over study in healthy middle-aged subjects.](#)  
Nilsson AC, Johansson-Boll EV, Björck IM.  
Br J Nutr. 2015 Sep 28;114(6):899-907. doi: 10.1017/S0007114515002524. Epub 2015 Aug 11.  
PMID: 26259632 Clinical Trial.
46. [Assessment of Blood Glucose Regulation and Safety of Resistant Starch Formula-Based Diet in Healthy Normal and Subjects With Type 2 Diabetes.](#)  
Lin CH, Chang DM, Wu DJ, Peng HY, Chuang LM.  
Medicine (Baltimore). 2015 Aug;94(33):e1332. doi: 10.1097/MD.0000000000001332.  
PMID: 26287417 **Free PMC article.** Clinical Trial.
47. [Effects of Acute Ingestion of Native Banana Starch on Glycemic Response Evaluated by Continuous Glucose Monitoring in Obese and Lean Subjects.](#)

- Jiménez-Domínguez G, Ble-Castillo JL, Aparicio-Trápala MA, Juárez-Rojop IE, Tovilla-Zárate CA, Ble-Castillo DJ, García-Vázquez C, Olvera-Hernández V, Pérez-Pimienta B, Diaz-Zagoya JC, Mendez JD. *Int J Environ Res Public Health.* 2015 Jul 6;12(7):7491-505. doi: 10.3390/ijerph120707491. PMID: 26154657 **Free PMC article.** Clinical Trial.
48. [High Amylose White Rice Reduces Post-Prandial Glycemic Response but Not Appetite in Humans.](#)  
Zenel AM, Stewart ML.  
*Nutrients.* 2015 Jul 2;7(7):5362-74. doi: 10.3390/nu7075225.  
PMID: 26147654 **Free PMC article.** Clinical Trial.
49. [Bioequivalence of n-3 fatty acids from microencapsulated fish oil formulations in human subjects.](#)  
Sanguansri L, Augustin MA, Lockett TJ, Abeywardena MY, Royle PJ, Mano MT, Patten GS.  
*Br J Nutr.* 2015 Mar 14;113(5):822-31. doi: 10.1017/S000711451400436X. Epub 2015 Feb 25.  
PMID: 25711158 Clinical Trial.
50. [Effect of cooling of cooked white rice on resistant starch content and glycemic response.](#)  
Sonia S, Witjaksono F, Ridwan R.  
*Asia Pac J Clin Nutr.* 2015;24(4):620-5. doi: 10.6133/apjcn.2015.24.4.13.  
PMID: 26693746 Clinical Trial.
51. [Resistant starch lowers postprandial glucose and leptin in overweight adults consuming a moderate-to-high-fat diet: a randomized-controlled trial.](#)  
Maziarz MP, Preisendanz S, Juma S, Imrhan V, Prasad C, Vijayagopal P.  
*Nutr J.* 2017 Feb 21;16(1):14. doi: 10.1186/s12937-017-0235-8.  
PMID: 28222742 **Free PMC article.** Clinical Trial.
52. [The acute effects of inulin and resistant starch on postprandial serum short-chain fatty acids and second-meal glycemic response in lean and overweight humans.](#)  
Rahat-Rozenbloom S, Fernandes J, Cheng J, Gloor GB, Wolever TM.  
*Eur J Clin Nutr.* 2017 Feb;71(2):227-233. doi: 10.1038/ejcn.2016.248. Epub 2016 Dec 14.  
PMID: 27966565 **Free PMC article.** Clinical Trial.
53. [Diets high in resistant starch increase plasma levels of trimethylamine-N-oxide, a gut microbiome metabolite associated with CVD risk.](#)  
Bergeron N, Williams PT, Lamendella R, Faghihnia N, Grube A, Li X, Wang Z, Knight R, Jansson JK, Hazen SL, Krauss RM.  
*Br J Nutr.* 2016 Dec;116(12):2020-2029. doi: 10.1017/S0007114516004165. Epub 2016 Dec 20.  
PMID: 27993177 **Free PMC article.** Clinical Trial.
54. [Resistant Starch Bagels Reduce Fasting and Postprandial Insulin in Adults at Risk of Type 2 Diabetes.](#)  
Dainty SA, Klingel SL, Pilkey SE, McDonald E, McKeown B, Emes MJ, Duncan AM.  
*J Nutr.* 2016 Nov;146(11):2252-2259. doi: 10.3945/jn.116.239418. Epub 2016 Oct 12.  
PMID: 27733521 Clinical Trial.
55. [Reformulating cereal bars: high resistant starch reduces in vitro digestibility but not in vivo glucose or insulin response; whey protein reduces glucose but disproportionately increases insulin.](#)  
Wolever TM, van Klinken BJ, Bordenave N, Kaczmarczyk M, Jenkins AL, Chu Y, Harkness L.  
*Am J Clin Nutr.* 2016 Oct;104(4):995-1003. doi: 10.3945/ajcn.116.132431. Epub 2016 Aug 31.  
PMID: 27581470 Clinical Trial.

56. [Effects of wheat bran extract rich in arabinoxylan oligosaccharides and resistant starch on overnight glucose tolerance and markers of gut fermentation in healthy young adults.](#)  
Boll EV, Ekström LM, Courtin CM, Delcour JA, Nilsson AC, Björck IM, Östman EM.  
Eur J Nutr. 2016 Jun;55(4):1661-70. doi: 10.1007/s00394-015-0985-z. Epub 2015 Jul 14.  
PMID: 26169871 Clinical Trial.
57. [Including dietary fiber and resistant starch to increase satiety and reduce aggression in gestating sows.](#)  
Sapkota A, Marchant-Forde JN, Richert BT, Lay DC.  
J Anim Sci. 2016 May;94(5):2117-27. doi: 10.2527/jas.2015-0013.  
PMID: 27285708 Clinical Trial.
58. [A novel fiber composite ingredient incorporated into a beverage and bar blunts postprandial serum glucose and insulin responses: a randomized controlled trial.](#)  
O'Connor LE, Campbell WW.  
Nutr Res. 2016 Mar;36(3):253-61. doi: 10.1016/j.nutres.2015.11.015. Epub 2015 Nov 26.  
PMID: 26923512 Clinical Trial.
59. [Glycemic Index of a Novel High-Fiber White Rice Variety Developed in India--A Randomized Control Trial Study.](#)  
Mohan V, Anjana RM, Gayathri R, Ramya Bai M, Lakshmipriya N, Ruchi V,  
Balasubramaniyam KK, Jakir MM, Shobana S, Unnikrishnan R, Krishnaswamy K, Henry JK, Sudha V.  
Diabetes Technol Ther. 2016 Mar;18(3):164-70. doi: 10.1089/dia.2015.0313. Epub 2016 Jan 7.  
PMID: 26741823 Clinical Trial.
60. [Effects of total fibre or resistant starch-rich diets within lifestyle intervention in obese prediabetic adults.](#)  
Dodevska MS, Sobajic SS, Djordjevic PB, Dimitrijevic-Sreckovic VS, Spasojevic-Kalimanovska VV, Djordjevic BI.  
Eur J Nutr. 2016 Feb;55(1):127-37. doi: 10.1007/s00394-015-0831-3. Epub 2015 Jan 15.  
PMID: 25588971 Clinical Trial.
61. [Postprandial glucose and insulin response to a high-fiber muffin top containing resistant starch type 4 in healthy adults: a double-blind, randomized, controlled trial.](#)  
Stewart ML, Zimmer JP.  
Nutrition. 2018 Sep;53:59-63. doi: 10.1016/j.nut.2018.01.002. Epub 2018 Feb 13.  
PMID: 29655779 Clinical Trial.
62. [Role of resistant starch on diabetes risk factors in people with prediabetes: Design, conduct, and baseline results of the STARCH trial.](#)  
Marlatt KL, White UA, Beyl RA, Peterson CM, Martin CK, Marco ML, Keenan MJ, Martin RJ, Aryana KJ, Ravussin E.  
Contemp Clin Trials. 2018 Feb;65:99-108. doi: 10.1016/j.cct.2017.12.005. Epub 2017 Dec 21.  
PMID: 29274892 **Free PMC article.** Clinical Trial.
63. [Type-4 Resistant Starch in Substitution for Available Carbohydrate Reduces Postprandial Glycemic Response and Hunger in Acute, Randomized, Double-Blind, Controlled Study.](#)  
Stewart ML, Wilcox ML, Bell M, Buggia MA, Maki KC.  
Nutrients. 2018 Jan 26;10(2):129. doi: 10.3390/nu10020129.  
PMID: 29373530 **Free PMC article.** Clinical Trial.
64. [Effect of Preexercise Ingestion of Modified Amylomaize Starch on Glycemic Response While Cycling.](#)  
Parks RB, Angus HF, King DS, Sharp RL.  
Int J Sport Nutr Exerc Metab. 2018 Jan 1;28(1):82-89. doi: 10.1123/ijsnem.2017-0193.  
Epub 2018 Jan 22.

- PMID: 29035602 Clinical Trial.
65. [Effect of resistant wheat starch on subjective appetite and food intake in healthy adults.](#)  
Emilien CH, Hsu WH, Hollis JH.  
*Nutrition*. 2017 Nov-Dec;43-44:69-74. doi: 10.1016/j.nut.2017.06.020. Epub 2017 Jul 6.  
PMID: 28935147 Clinical Trial.
66. [Investigating the relationship between lentil carbohydrate fractions and in vivo postprandial blood glucose response by use of the natural variation in starch fractions among 20 lentil varieties.](#)  
Ramdath DD, Liu Q, Donner E, Hawke A, Kalinga D, Winberg J, Wolever TMS.  
*Food Funct.* 2017 Oct 18;8(10):3783-3791. doi: 10.1039/c7fo00972k.  
PMID: 28959813 Clinical Trial.
67. [Acute increases in serum colonic short-chain fatty acids elicited by inulin do not increase GLP-1 or PYY responses but may reduce ghrelin in lean and overweight humans.](#)  
Rahat-Rozenbloom S, Fernandes J, Cheng J, Wolever TMS.  
*Eur J Clin Nutr.* 2017 Aug;71(8):953-958. doi: 10.1038/ejcn.2016.249. Epub 2016 Dec 14.  
PMID: 27966574 **Free PMC article.** Clinical Trial.
68. [Effects of whole grain rye, with and without resistant starch type 2 supplementation, on glucose tolerance, gut hormones, inflammation and appetite regulation in an 11-14.5 hour perspective; a randomized controlled study in healthy subjects.](#)  
Sandberg JC, Björck IME, Nilsson AC.  
*Nutr J.* 2017 Apr 21;16(1):25. doi: 10.1186/s12937-017-0246-5.  
PMID: 28431559 **Free PMC article.** Clinical Trial.
69. [A High Fiber Cookie Made with Resistant Starch Type 4 Reduces Post-Prandial Glucose and Insulin Responses in Healthy Adults.](#)  
Stewart ML, Zimmer JP.  
*Nutrients.* 2017 Mar 5;9(3):237. doi: 10.3390/nu9030237.  
PMID: 28273870 **Free PMC article.** Clinical Trial.
70. [Chilled Potatoes Decrease Postprandial Glucose, Insulin, and Glucose-dependent Insulinotropic Peptide Compared to Boiled Potatoes in Females with Elevated Fasting Glucose and Insulin.](#)  
Patterson MA, Fong JN, Maiya M, Kung S, Sarkissian A, Nashef N, Wang W.  
*Nutrients.* 2019 Sep 3;11(9):2066. doi: 10.3390/nu11092066.  
PMID: 31484331 **Free PMC article.** Clinical Trial.
71. [High-Resistant Starch, Low-Protein Flour Intervention on Patients With Early Type 2 Diabetic Nephropathy: A Randomized Trial.](#)  
Meng Y, Bai H, Yu Q, Yan J, Zhao L, Wang S, Li Z, Wang Q, Chen L.  
*J Ren Nutr.* 2019 Sep;29(5):386-393. doi: 10.1053/j.jrn.2018.12.005. Epub 2019 Apr 11.  
PMID: 30982743 Clinical Trial.
72. [High-Amylose Wheat Lowers the Postprandial Glycemic Response to Bread in Healthy Adults: A Randomized Controlled Crossover Trial.](#)  
Belobrajdic DP, Regina A, Klingner B, Zajac I, Chapron S, Berbezy P, Bird AR.  
*J Nutr.* 2019 Aug 1;149(8):1335-1345. doi: 10.1093/jn/nxz067.  
PMID: 31162585 Clinical Trial.
73. [Beneficial effects of green banana biomass consumption in patients with pre-diabetes and type 2 diabetes: a randomised controlled trial.](#)  
Costa ES, França CN, Fonseca FAH, Kato JT, Bianco HT, Freitas TT, Fonseca HAR, Figueiredo Neto AM, Izar MC.  
*Br J Nutr.* 2019 Jun;121(12):1365-1375. doi: 10.1017/S0007114519000576. Epub 2019 Mar 19.  
PMID: 30887937 Clinical Trial.

74. [Potato product form impacts in vitro starch digestibility and glucose transport but only modestly impacts 24 h blood glucose response in humans.](#)  
Li M , George J , Hunter S , Hamaker B , Mattes R , Ferruzzi MG .  
Food Funct. 2019 Apr 17;10(4):1846-1855. doi: 10.1039/c8fo02530d.  
PMID: 30942789 Clinical Trial.
75. [Metabolic phenotypes and the gut microbiota in response to dietary resistant starch type 2 in normal-weight subjects: a randomized crossover trial.](#)  
Zhang L, Ouyang Y, Li H, Shen L, Ni Y, Fang Q, Wu G, Qian L, Xiao Y, Zhang J, Yin P, Panagiotou G, Xu G, Ye J, Jia W.  
Sci Rep. 2019 Mar 20;9(1):4736. doi: 10.1038/s41598-018-38216-9.  
PMID: 30894560 **Free PMC article.** Clinical Trial.
76. [Acute Effects of Three Cooked Non-Cereal Starchy Foods on Postprandial Glycemic Responses and in Vitro Carbohydrate Digestion in Comparison with Whole Grains: A Randomized Trial.](#)  
Zhu R, Fan Z, Han Y, Li S, Li G, Wang L, Ye T, Zhao W.  
Nutrients. 2019 Mar 15;11(3):634. doi: 10.3390/nu11030634.  
PMID: 30875961 **Free PMC article.** Clinical Trial.
77. [Acute Effect of Resistant Starch on Food Intake, Appetite and Satiety in Overweight/Obese Males.](#)  
Al-Mana NM, Robertson MD.  
Nutrients. 2018 Dec 15;10(12):1993. doi: 10.3390/nu10121993.  
PMID: 30558330 **Free PMC article.** Clinical Trial.
78. [Impact of rye-based evening meals on cognitive functions, mood and cardiometabolic risk factors: a randomized controlled study in healthy middle-aged subjects.](#)  
Sandberg JC, Björck IME, Nilsson AC.  
Nutr J. 2018 Nov 6;17(1):102. doi: 10.1186/s12937-018-0412-4.  
PMID: 30400947 **Free PMC article.** Clinical Trial.
79. [Effect of 12 wk of resistant starch supplementation on cardiometabolic risk factors in adults with prediabetes: a randomized controlled trial.](#)  
Peterson CM, Beyl RA, Marlatt KL, Martin CK, Aryana KJ, Marco ML, Martin RJ, Keenan MJ, Ravussin E.  
Am J Clin Nutr. 2018 Sep 1;108(3):492-501. doi: 10.1093/ajcn/nqy121.  
PMID: 30010698 **Free PMC article.** Clinical Trial.
80. [Resistant Starch Type 2 from Wheat Reduces Postprandial Glycemic Response with Concurrent Alterations in Gut Microbiota Composition.](#)  
Hughes RL, Horn WH, Finnegan P, Newman JW, Marco ML, Keim NL, Kable ME.  
Nutrients. 2021 Feb 17;13(2):645. doi: 10.3390/nu13020645.  
PMID: 33671147 **Free PMC article.** Clinical Trial.
81. [Noodles Made from High Amylose Wheat Flour Attenuate Postprandial Glycaemia in Healthy Adults.](#)  
Ang K, Bourgy C, Fenton H, Regina A, Newberry M, Diepeveen D, Lafiandra D, Grafenauer S, Hunt W, Solah V.  
Nutrients. 2020 Jul 22;12(8):2171. doi: 10.3390/nu12082171.  
PMID: 32707905 **Free PMC article.** Clinical Trial.
82. [Nondigestible Carbohydrates Affect Metabolic Health and Gut Microbiota in Overweight Adults after Weight Loss.](#)  
Johnstone AM, Kelly J, Ryan S, Romero-Gonzalez R, McKinnon H, Fyfe C, Naslund E, Lopez-Nicolas R, Bosscher D, Bonnema A, Frontela-Saseta C, Ros-Berruezo G, Horgan G, Ze X, Harrold J, Halford J, Gratz SW, Duncan SH, Shirazi-Beechey S, Flint HJ.  
J Nutr. 2020 Jul 1;150(7):1859-1870. doi: 10.1093/jn/nxaa124.  
PMID: 32510158 Clinical Trial.

83. [Food prototype containing resistant starch type 4 on postprandial glycemic response in healthy adults.](#)  
Du Y, Wu Y, Xiao D, Guzman G, Stewart ML, Gourineni V, Burton-Freeman B, Edirisinghe I.  
Food Funct. 2020 Mar 26;11(3):2231-2237. doi: 10.1039/c9fo02674f.  
PMID: 32101207 Clinical Trial.
84. [Effects of single ingestion of rice cracker and cooked rice with high resistant starch on postprandial glucose and insulin responses in healthy adults: two randomized, single-blind, cross-over trials.](#)  
Saito Y, Watanabe T, Sasaki T, Watanabe K, Hirayama M, Fujita N.  
Biosci Biotechnol Biochem. 2020 Feb;84(2):365-371. doi:  
10.1080/09168451.2019.1687282. Epub 2019 Nov 5.  
PMID: 31690219 Clinical Trial.
85. [The In Vivo Net Energy Content of Resistant Starch and Its Effect on Macronutrient Oxidation in Healthy Adults.](#)  
Giles ED, Brown IL, MacLean PS, Pan Z, Melanson EL, Heard KJ, Cornier MA, Marden T, Higgins JA.  
Nutrients. 2019 Oct 16;11(10):2484. doi: 10.3390/nu11102484.  
PMID: 31623184 **Free PMC article.** Clinical Trial.

## Blood Sugar Management: Oat Beta-Glucan Studies (19)

1. [Effect of processing on oat β-glucan viscosity, postprandial glycemic response and subjective measures of appetite.](#)  
Ames N , Malunga LN , Mollard R , Johnson J , Chu Y , Thandapilly SJ .  
Food Funct. 2021 Apr 21;12(8):3672-3679. doi: 10.1039/d0fo03283b. Epub 2021 Apr 1.  
PMID: 33900322 Clinical Trial.
2. [Effect of Varying Molecular Weight of Oat β-Glucan Taken just before Eating on Postprandial Glycemic Response in Healthy Humans.](#)  
Wolever TMS, Mattila O, Rosa-Sibakov N, Tosh SM, Jenkins AL, Ezatagha A, Duss R, Steinert RE.  
Nutrients. 2020 Jul 29;12(8):2275. doi: 10.3390/nu12082275.  
PMID: 32751269 **Free PMC article.** Clinical Trial.
3. [Increasing oat β-glucan viscosity in a breakfast meal slows gastric emptying and reduces glycemic and insulinemic responses but has no effect on appetite, food intake, or plasma ghrelin and PYY responses in healthy humans: a randomized, placebo-controlled, crossover trial.](#)  
Wolever TMS, Tosh SM, Spruill SE, Jenkins AL, Ezatagha A, Duss R, Johnson J, Chu Y, Steinert RE.  
Am J Clin Nutr. 2020 Feb 1;111(2):319-328. doi: 10.1093/ajcn/nqz285.

- PMID: 31828287 Clinical Trial.
4. [Impact of oat processing on glycaemic and insulinaemic responses in healthy humans: a randomised clinical trial.](#)  
Wolever TMS, Johnson J, Jenkins AL, Campbell JC, Ezatagha A, Chu Y.  
Br J Nutr. 2019 Jun;121(11):1264-1270. doi: 10.1017/S0007114519000370. Epub 2019 May 9.  
PMID: 31068229 Clinical Trial.
5. [Effects of oat  \$\beta\$ -glucan consumption at breakfast on ad libitum eating, appetite, glycemia, insulinemia and GLP-1 concentrations in healthy subjects.](#)  
Zaremba SMM, Gow IF, Drummond S, McCluskey JT, Steinert RE.  
Appetite. 2018 Sep 1;128:197-204. doi: 10.1016/j.appet.2018.06.019. Epub 2018 Jun 18.  
PMID: 29920323 Clinical Trial.
6. [Effect of adding oat bran to instant oatmeal on glycaemic response in humans - a study to establish the minimum effective dose of oat  \$\beta\$ -glucan.](#)  
Wolever TMS, Jenkins AL, Prudence K, Johnson J, Duss R, Chu Y, Steinert RE.  
Food Funct. 2018 Mar 1;9(3):1692-1700. doi: 10.1039/c7fo01768e. Epub 2018 Feb 26.  
PMID: 29480316 Clinical Trial.
7. [Effect of Consuming Oat Bran Mixed in Water before a Meal on Glycemic Responses in Healthy Humans-A Pilot Study.](#)  
Steinert RE, Raederstorff D, Wolever TM.  
Nutrients. 2016 Aug 26;8(9):524. doi: 10.3390/nu8090524.  
PMID: 27571099 **Free PMC article.** Clinical Trial.
8. [Increasing the viscosity of oat  \$\beta\$ -glucan beverages by reducing solution volume does not reduce glycaemic responses.](#)  
Kwong MG, Wolever TM, Brummer Y, Tosh SM.  
Br J Nutr. 2013 Oct;110(8):1465-71. doi: 10.1017/S000711451300069X. Epub 2013 Jun 21.  
PMID: 23789885 Clinical Trial.
9. [Semisolid meal enriched in oat bran decreases plasma glucose and insulin levels, but does not change gastrointestinal peptide responses or short-term appetite in healthy subjects.](#)  
Juvonen KR, Salmenkallio-Marttila M, Lylly M, Liukkonen KH, Lähteenmäki L, Laaksonen DE, Uusitupa MI, Herzig KH, Poutanen KS, Karhunen LJ.  
Nutr Metab Cardiovasc Dis. 2011 Sep;21(9):748-56. doi: 10.1016/j.numecd.2010.02.002. Epub 2010 Jun 4.

- PMID: 20605427 Clinical Trial.
10. [beta-Glucan-enriched bread reduces energy intake and modifies plasma ghrelin and peptide YY concentrations in the short term.](#)  
Vitaglione P, Lumaga RB, Stanzone A, Scalfi L, Fogliano V.  
Appetite. 2009 Dec;53(3):338-44. doi: 10.1016/j.appet.2009.07.013. Epub 2009 Jul 23.  
PMID: 19631705 Clinical Trial.
11. [Physicochemical properties of beta-glucan in differently processed oat foods influence glycemic response.](#)  
Regand A, Tosh SM, Wolever TM, Wood PJ.  
J Agric Food Chem. 2009 Oct 14;57(19):8831-8. doi: 10.1021/jf901271v.  
PMID: 19728711 Clinical Trial.
12. [Viscosity of oat bran-enriched beverages influences gastrointestinal hormonal responses in healthy humans.](#)  
Juvonen KR, Purhonen AK, Salmenkallio-Marttila M, Lähteenmäki L, Laaksonen DE, Herzig KH, Uusitupa MI, Poutanen KS, Karhunen LJ.  
J Nutr. 2009 Mar;139(3):461-6. doi: 10.3945/jn.108.099945. Epub 2009 Jan 28.  
PMID: 19176745 Clinical Trial.
13. [Effect of muesli with 4 g oat beta-glucan on postprandial blood glucose, gastric emptying and satiety in healthy subjects: a randomized crossover trial.](#)  
Hlebowicz J, Darwiche G, Björnell O, Almér LO.  
J Am Coll Nutr. 2008 Aug;27(4):470-5. doi: 10.1080/07315724.2008.10719727.  
PMID: 18978166 Clinical Trial.
14. [Serum lipids and postprandial glucose and insulin levels in hyperlipidemic subjects after consumption of an oat beta-glucan-containing ready meal.](#)  
Biörklund M, Holm J, Onning G.  
Ann Nutr Metab. 2008;52(2):83-90. doi: 10.1159/000121281. Epub 2008 Mar 11.  
PMID: 18334815 Clinical Trial.
15. [Beta-glucan from two sources of oat concentrates affect postprandial glycemia in relation to the level of viscosity.](#)  
Panahi S, Ezatagha A, Temelli F, Vasanthan T, Vuksan V.  
J Am Coll Nutr. 2007 Dec;26(6):639-44. doi: 10.1080/07315724.2007.10719641.  
PMID: 18187427 Clinical Trial.

16. [Effects of consuming foods containing oat beta-glucan on blood pressure, carbohydrate metabolism and biomarkers of oxidative stress in men and women with elevated blood pressure.](#)

Maki KC, Galant R, Samuel P, Tesser J, Witchger MS, Ribaya-Mercado JD, Blumberg JB, Geohas J.

Eur J Clin Nutr. 2007 Jun;61(6):786-95. doi: 10.1038/sj.ejcn.1602562. Epub 2006 Dec 6.

PMID: 17151592 Clinical Trial.

17. [Concentrated oat beta-glucan, a fermentable fiber, lowers serum cholesterol in hypercholesterolemic adults in a randomized controlled trial.](#)

Queenan KM, Stewart ML, Smith KN, Thomas W, Fulcher RG, Slavin JL.

Nutr J. 2007 Mar 26;6:6. doi: 10.1186/1475-2891-6-6.

PMID: 17386092 **Free PMC article.** Clinical Trial.

18. [Sweeteners and beta-glucans improve metabolic and anthropometrics variables in well controlled type 2 diabetic patients.](#)

Reyna NY, Cano C, Bermúdez VJ, Medina MT, Souki AJ, Ambard M, Nuñez M, Ferrer MA, Inglett GE.

Am J Ther. 2003 Nov-Dec;10(6):438-43. doi: 10.1097/00045391-200311000-00010.

PMID: 14624282 Clinical Trial.

19. [Postprandial glucose, insulin, and incretin responses to grain products in healthy subjects.](#)

Juntunen KS, Niskanen LK, Liukkonen KH, Poutanen KS, Holst JJ, Mykkänen HM.

Am J Clin Nutr. 2002 Feb;75(2):254-62. doi: 10.1093/ajcn/75.2.254.

PMID: 11815315 Clinical Trial.